

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/14, 9/00	A1	(11) International Publication Number: WO 96/20696 (43) International Publication Date: 11 July 1996 (11.07.96)
(21) International Application Number: PCT/US95/16445 (22) International Filing Date: 4 December 1995 (04.12.95) (30) Priority Data: 08/366,518 30 December 1994 (30.12.94) US (71) Applicant (for all designated States except US): EASTMAN KODAK COMPANY [US/US]; 343 State Street, Rochester, NY 14650 (US). (72) Inventors; and (73) Inventors/Applicants (for US only): RUDDY, Stephen, B. [US/US]; 226 Stallion Lane, Schwenksville, PA 19473 (US); EICKHOFF, W., Mark [US/US]; 1313 Rhode Island Circle, Downingtown, PA 19355 (US); LIVERSIDGE, Gary [US/US]; 258 Colwyn Terrace, West Chester, PA 19380 (US); COOPER, Eugene, R. [US/US]; 2612 Crum Creek Drive, Berwyn, PA 19312 (US). (74) Agent: ROSENSTEIN, Arthur, H.; 343 State Street, Rochester, NY 14650-2201 (US).	(81) Designated States: AL, AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LS, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: FORMULATIONS OF ORAL GASTROINTESTINAL THERAPEUTIC AGENTS IN COMBINATION WITH PHARMACEUTICALLY ACCEPTABLE CLAYS (57) Abstract Nanoparticulate crystalline therapeutic substances formulated with stabilizers and pharmaceutically acceptable clays to enhance contact between the crystalline therapeutic substances and the gastrointestinal tract and to provide extended therapeutic effect.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MF	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Ghana	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BT	Bhutan	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KZ	Republic of Korea	SE	Sweden
CG	Congo	KM	Komoros	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SE	Switzerland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	MH	Marshall Islands	UG	Uganda
FI	Finland	ML	Mali	US	United States of America
FR	France	MN	Mongolia	UZ	Uzbekistan
GA	Gabon	MR	Mauritania	VN	Viet Nam

Field of the Invention

10

15

25

30

35

Bioadhesion is usually achieved by interaction of either a synthetic or natural polymeric substance with the mucosal membranes of the GI tract. Such technology has been employed to enhance drug delivery by decreasing the transit time of a drug substance in the GI tract and hence promote an opportunity for enhanced absorption. With regards to the development water-insoluble or poorly water-soluble drug formulations intended to coat the GI tract, it is important to identify mucosal adhesives that coat the GI surfaces and affect diseased or abnormal tissues. Highly charged carboxylated polyanions are good candidates for

use as bioadhesives in the GI tract. See, for example: Park, K. and Robinson, J.R., Bioadhesion: Polymers and Platforms for Oral-Controlled Drug Delivery; Method to Study Bioadhesion. *Int. J. Pharm.*, 19, 107 (1984). The formation of a bioadhesive bond between a polymeric substance and the mucosal lining of the GI tract can be visualized as a two step process, i.e., initial contact between the two surfaces and the formation of secondary bonds due to non-covalent interactions. Bioadhesives specific for the GI tract must interact with the mucus layer during attachment. Mucus, a general term for the heterogenous secretion found on the epithelial surfaces of the GI tract, is made of the following components:

glycoprotein macromolecules, inorganic salts, proteins, lipids and mucopolysaccharides. These glycoproteins typically consist of a protein core with carbohydrate side chains. This forms a network of mucus that is a continuous layer covering the GI tract. From a bioadhesive perspective, mucus consists of highly hydrated, crosslinked linear, flexible yet random coiled glycoprotein molecules with a net negative charge. Understanding the principles of bioadhesion is the basis for formulating an oral compositions for coating the GI tract. Bioadhesion accounts for the interaction between a biological surface and a biomaterial substance. As noted previously, bioadhesive agents are usually polymeric substances that adhere to tissues by ionic or covalent bonds or by physical attachment. Several theories of bioadhesion have been published including electronic, adsorption, wetting, diffusion and fracture theories. Bioadhesives bind to membrane surfaces and are retained for variable periods of time.

We have now discovered that certain therapeutic drugs in a combination with bioadhesive or mucoadhesive surfactants and pharmaceutically acceptable clays provide excellent coating on the GI tract for a prolonged period of time so that the therapeutic drugs are able to affect diseased conditions which may be present in the GI tract.

In accordance with the present invention, there is provided an orally/rectally administrable GI formulations containing an effective amount of a water-insoluble or poorly water-soluble therapeutic agent. There is further provided a method for affecting diseased conditions in the GI tract comprising oral or rectal administration to a patient an effective amount of the above-identified formulation to prevent or cure such diseased conditions.

Summary Of The Invention

In accordance with the present invention there is provided an orally/rectally administrable therapeutic composition comprising:

5 of from about 4 to about 45% w/v, and preferably of from about 5 to about 25% w/v, of an essentially water insoluble or poorly water-soluble particulate crystalline material having an effective average particle size of less than about 2,000 nm, more preferably an effective average particle size of less than about 1,000 nm, and most preferably an effective average particle size of less than about 400 nm;

10 of from about 0.1% to about 10% w/v, and preferably of from about 1% to about 6% w/v of a bioadhesive or mucoadhesive surfactant stabilizer (hereinafter sometimes referred to as "primary stabilizer");

15 of from about 0.1 to about 10% w/v and preferably of from about 0.5 to about 5, and most preferably of from about 1 to 2% of a pharmaceutically acceptable clay; and

water to make 100% w/v.

20 Secondary stabilizers may also be used in the x-ray contrast formulation up to about 1% w/v, preferably up to about 0.2% w/v, and most preferably up to about 0.1% w/v. Secondary stabilizers include dioctylsulfosuccinate (DOSS) and sodium lauryl sulfate (SLS).

Other ingredients customarily used in oral pharmaceutical formulations may also be included, such as flavorants, colorants and preservatives to provide pharmaceutically acceptable and palatable formulations.

25 Detailed Description Of The Invention

The present invention is based on the discovery that particulate crystalline materials can be rendered mucoadhesive or bioadhesive in the gastrointestinal tract when the particulate crystalline material is combined with certain surface active agents in a suspension.

30 The invention can be practiced with a wide variety of crystalline materials that are water-insoluble or poorly soluble in water. As used herein "poorly soluble" means that the material has a solubility in aqueous medium of less than about 10 mg/ml, and preferably of less than about 1 mg/ml. Suitable drug substances for use in the present invention follow.

Drugs

- Suitable drug substance can be selected from a variety of known classes of drugs including, for example, antacids, anti-inflammatory agents, antibiotics (including penicillins), antimycobacterial agents, antiviral agents, corticosteroids, 5 parasympathomimetics, radio-pharmaceuticals, sympathomimetics, demulcents, emollients, gastrointestinal protectives and adsorbents, antifungals, H₂-blocking agents, proton pump inhibitors, muscarinic antagonists, bismuth compounds, sucralfate, carbenoxolone, prostaglandins, digestants, bile acids, laxatives, antiparasitic agents, anthelmintics, antiprotozoal agents, antimicrobial agents, 10 vitamins, immunologic agents, vaccines, anesthetics, lipid-regulating agents and bile acid sequestrants. Preferred drug substances include those intended for oral administration and rectal administration. A description of these classes of drugs and a listing of species within each class can be found in Martindale, The Extra Pharmacopoeia, Twenty-Ninth Edition, The Pharmaceutical Press, London, 1989, 15 the disclosure of which is hereby incorporated by reference in its entirety. The drug substances are commercially available and/or can be prepared by techniques known in the art.

Bioadhesives/Mucoadhesives

- 20 The surfactants found to have bioadhesive or mucoadhesive properties include:

- 1) Poloxamers having an average molecular weight of from about 1,000 to 15,000 daltons;
- 2) Polyvinyl alcohol;
- 25 3) Polyvinyl pyrrolidone,
- 4) Hydroxypropyl methylcellulose; and
- 5) Polyoxyethylene sorbitan mono-oleate (Tween 80).

- Poloxamers are polyethylene-polypropylene glycol block polymers containing ethylene oxide (PEO) and propylene oxide (PPO) moles according to 30 the formula (PEO)^a - (PPO)^b - (PEO)^c wherein

a is 46, 52, 62, 75, 97, 98, 122 and 128;

b is 16, 30, 35, 39, 47, 54 and 67; and

c is 46, 52, 62, 75, 97, 98, 122 and 128.

- 35 Table 1 shows the various poloxamers by manufacturer-designated number.

TABLE 1
Molecular Weights of Poloxamers

	Poloxamer No.	Pluronic	Av. Mol. Wt.	Av. Values		
				a	b	c
5	401		4,400	6	67	6
	402		5,000	13	67	13
	403		5,750	21	67	21
	407	F127	12,000	98	67	98
10	331		3,800	7	54	7
	333		4,950	20	54	20
	334		5,850	31	54	31
	335		6,000	38	54	38
	338	F108	15,000	128	54	128
15	282		3,650	10	47	10
	284		4,600	21	47	21
	288	F98	13,500	122	47	122
	231		2,750	6	39	6
	234		4,200	22	39	22
20	235		4,600	27	39	27
	237	F87	7,700	62	39	62
	238	F88	10,800	97	39	97
	212		2,750	8	35	8
	215		4,150	24	35	24
25	217	F77	6,600	52	35	52
	181		2,000	3	30	3
	182		2,500	8	30	8
	183		2,650	10	30	10
	184		2,900	13	30	13
30	185		3,400	19	30	19
	188	F68	8,350	75	30	75
	122		1,630	5	21	5
	123		1,850	7	21	7
	124		2,200	11	21	11
35	101		1,100	2	16	2
	105		1,900	11	16	11

108 F38 5,000 46 16 46

Certain number of these surfactants are also known as Pluronic, which is a brand name of BASF Corporation.

Preferred surfactants for use in the present invention are:

5

Pluronic F127

Pluronic F108

Pluronic F98

Pluronic F87

10

Pluronic F88

Pluronic 77

Pluronic F68 and

Pluronic F38.

15 Method of Preparing the Radiopaque Particulates

The particulates were prepared by milling the large particles mixed with an appropriate surface active agent to obtain the desired particle size. Alternatively, the large particulates may be comminuted to the desired particle size and subsequently intimately mixed with the appropriate surface active agent. The milling technique is described in U.S. Patent No. 5,145,684, which is incorporated herein by reference.

As used herein, particle size refers to a number average particle size as measured by conventional particle size measuring techniques well known to those skilled in the art, such as sedimentation field flow fractionation, photon correlation spectroscopy, or disk centrifugation. By "an effective average particle size of less than about 400 nm" for example, it is meant that at least 90% of the particles have a weight average particle size of less than about 400 nm when measured by the above-noted techniques. With reference to the effective average particle size, it is preferred that at least 95% and, more preferably, at least 99% of the particles have a particle size less than the effective average, e.g., 400 nm. In particularly preferred embodiments, essentially all of the particles have a size less than 400 nm.

The particles of this invention can be prepared in a method comprising the steps of dispersing a substance in a liquid dispersion medium and applying mechanical means in the presence of grinding media to reduce the particle size of the substance to an effective average particle size of less than about 400 nm. The particles are reduced in size in the presence of the surface active agent.

Alternatively, the particles can be intimately mixed with a surface active agent after attrition.

A general procedure for preparing the particles of this invention is set forth below. The substance selected is obtained commercially and/or prepared by techniques known in the art in a conventional coarse form. It is preferred, but not essential, that the particle size of the coarse substance selected be less than about 100 mm as determined by sieve analysis. If the coarse particle size of the substance is greater than about 100 mm, then it is preferred that the particles of the substance be reduced in size to less than 100 mm using a conventional milling method such as airjet or fragmentation milling.

The mechanical means applied to reduce the particle size of the substance conveniently can take the form of a dispersion mill. Suitable dispersion mills include a ball mill, an attritor mill, a vibratory mill, and media mills such as a sand mill and a bead mill. A media mill is preferred due to the relatively shorter milling time required to provide the intended result, i.e., the desired reduction in particle size.

The grinding media for the particle size reduction step can be selected from rigid media preferably spherical or particulate in form having an average size less than about 3 mm and, more preferably, less than about 1 mm. Such media desirably can provide the particles of the invention with shorter processing times and impart less wear to the milling equipment. The selection of material for the grinding media is not believed to be critical. We have found that zirconium oxide, such as 95% ZrO stabilized with magnesia, zirconium silicate and glass grinding media provide particles having levels of contamination which are believed to be acceptable for the preparation of pharmaceutical compositions. However, other media, such as stainless steel, titania, alumina, and 95% ZrO stabilized with yttrium, are expected to be useful. Preferred media have a density greater than about 3 g/cm³.

The attrition time can vary widely and depends primarily upon the particular mechanical means and processing conditions selected. For ball mills, processing times of up to five days or longer may be required. On the other hand, processing times of less than 1 day (residence times of one minute up to several hours) have provided the desired results using a high shear media mill.

The particles must be reduced in size at a temperature which does not significantly degrade the substance. Processing temperatures of less than about 30 °- 40°C are ordinarily preferred. If desired, the processing equipment can be

cooled with conventional cooling equipment. The method is conveniently carried out under conditions of ambient temperature and at processing pressures which are safe and effective for the milling process. For example, ambient processing pressures are typical of ball mills, attritor mills and vibratory mills. Processing pressures up to about 20 psi (1.4 kg/cm²) are typical of media milling.

Particle size analysis was carried out using the Microtract Ultrafine Particle Analyzer. (Leeds and Northrup Co.; St. Petersburg, Florida) Nanosuspension particle size was determined during the milling process and again immediately before the nanosuspensions were administered to rodents. Particle size was determined on the Coulter Model N4MD Submicron Particle Analyzer. (Coulter Corp.; Miami Lakes, Florida).

Formulation Examples

The following formulation examples will further illustrate the present invention.

Example 1

	Sucralfate	20 g
	Pluronic F127	4.0 g
20	Benzoate Sodium	0.2 g
	Saccharin Sodium	0.1 g
	FD&C Red No. 40	0.03 g
	Water, qs	100 ml

Example 2

25	Carbenoxolone	15 g
	Pluronic F127	4.0 g
	Benzoate Sodium	0.2 g
	Sorbate Potassium	0.15 g
30	Saccharin Sodium	0.1 g
	FD&C Red No. 40	0.03 g
	Water, qs	100 ml

Example 3

35	Ursodeoxycholic Acid	25 g
	Pluronic F88	5.0 g

-9-

	Benzoate Sodium	0.2 g
	Saccharin Sodium	0.1 g
	FD&C Red No. 3	0.03 g
	Water, qs	100 ml

5

Example 4

	Cholic Acid	19 g
	Sucrose	10 g
	Pluromic F77	4.0 g
10	Diocetylulfosuccinate	0.1 g
	Methylparabens	0.2 g
	Propylparabens	0.07 g
	FD&C Yellow No. 5	0.03 g
	Water, qs	100 ml

15

Example 5

	Vitamin B	15 g
	Pluronic F127	5 g
	Sorbitol	5 g
20	Benzoate Sodium	0.2 g
	Water, qs	100 ml

Example 6

	Penicillin	22 g
25	HPMC (2%=100 cps)	2 g
	Steam sterilized by autoclaving at 120_C for 21 minutes & 5 psig	0.2 g
	Water, qs	100 ml

Example 7

30	Antibacterial	20 g
	Pluronic F127	4.0 g
	Benzoate Sodium	0.2 g
	Sorbate Potassium	0.15 g
35	Saccharin Sodium	0.1 g
	Water, qs	100 ml

Hydrochloric Acid

adjust to pH 4.0

5 The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention, and all such modifications as would be obvious to one skilled in the art are intended to be included within the scope of the following claims.

WHAT IS CLAIMED IS:

1. An orally/rectally administrable therapeutic formulation comprising:
- 5 of from about 0.1 to about 45% w/v of an essentially water-insoluble particulate drug having an effective average particle size of less than about 2,000 nm;
- of from about 0.1 to about 10% w/v of a bioadhesive surfactant stabilizer
- 10 selected from the group consisting of: polyethylene-polypropylene glycol block polymers of the formula (i)
- (i) (polyethylene oxide)_a - (polypropylene oxide)_b - (polyethylene oxide)_c
- 15 wherein
- a is 46, 52, 62, 75, 97, 98, 122 and 128;
- b is 16, 30, 35, 39, 47, 54 and 67; and
- c is 46, 52, 62, 75, 97, 98, 122 and 128;
- (ii) polyvinyl alcohol,
- 20 (iii) polyvinyl pyrrolidone,
- (iv) hydroxypropyl methylcellulose, and
- (v) polyoxyethylene sorbitan mono-oleate;
- from about 0.1 to 10% of a pharmaceutically acceptable clay selected
- 25 from the group consisting of: montmorillonite, beidelite, nontronite, hectorite and saponite; and
- water to make 100% w/v.
- 30 2. The therapeutic formulation of claim 1 wherein said drug is selected from the group consisting of: antacids, anti-inflammatory agents, antibiotics, antimycobacterial agents, antiviral agents, corticosteroids, parasympathomimetics, radio-pharmaceuticals, sympathomimetics, demulcents, emollients, gastrointestinal protectives and adsorbents, antifungals, H₂-blocking agents, proton pump
- 35 inhibitors, muscarinic antagonists, bismuth compounds, sucralfate, carbenoxolone, prostaglandins, digestants, bile acids, laxatives, antiparasitic agents, anthelmintics,

antiprotozoal agents, antimicrobial agents, vitamins, immunologic agents, vaccines, anesthetics, lipid-regulating agents and bile acid sequestrants.

3. The therapeutic formulation of claim 1 wherein said pharmaceutically
5 acceptable clays is

montmorillonite, having the formula



- 10 beidelite, having the formula



nontronite, having the formula



- 15 wherein M^+ is Na, Ca or Mg.

4. The therapeutic formulation of claim 1 wherein said pharmaceutically
acceptable clays is

- 20 saponite, having the formula



hectorite, having the formula



- 25 wherein M^+ is Na, Ca or Mg.

INTERNATIONAL SEARCH REPORT

National Application No.

PCT/US 95/16445

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K9/14 A61K9/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Maximum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than maximum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP.A.0 331 385 (SATO PHARMACEUTICAL) 6 September 1989 see claims 1,2,4,5,8 see page 3, line 22 - line 32 -----	1,2

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is considered with one or more other such documents, such combination being obvious to a person skilled in the art.

B document member of the same patent family

Date of the actual completion of the international search

24 April 1996

Date of making of the international search report

- 6.05.96

Name and mailing address of the ISA

European Patent Office, P.O. Box 5115 Postfach 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Te. 31 651 490 41
Fax (+ 31-70) 340-2016

Authorized officer

Ventura Amat, A

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 95/16445

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-331385	06-09-89	JP-A- 1250323	05-10-89
		DE-D- 68914180	05-05-94
		DE-T- 68914180	11-08-94
		US-A- 5464828	07-11-95
